

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C12Q 1/58, 1/04</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/51769 (43) International Publication Date: 14 October 1999 (14.10.99)</p>
<p>(21) International Application Number: PCT/KR99/00161 (22) International Filing Date: 2 April 1999 (02.04.99) (30) Priority Data: 1998/11909 3 April 1998 (03.04.98) KR (71)(72) Applicants and Inventors: LEE, Jong-Hwa [KR/KR]; Yukyoung Apt., 105-109, Yobang-ri, Seongseo-cup, Cheonan-city, Choongcheongnam-do 330-830 (KR). LEE, Hak-Sung [KR/KR]; Deokpoong 1-dong, 420-52, Hanam-city, Kyungki-do 465-011 (KR). ROE, Im-Hwan [KR/KR]; Doosan Apt., 421-1401, Areum-maeul, Maesong-dong, Bundang-ku, Seongnam-city, Kyungki-do 463-065 (KR). KIM, Jung-Taik [KR/KR]; Hanyang Apt., 314-1601, Seohyeon-dong 91, Bundang-ku, Seongnam-city, Kyungki-do 463-050 (KR). HAH, Yung-Chil [KR/KR]; Hanyang Apt., 51-1201, Appoojeong-dong, Kangnam-ku, Seoul 135-110 (KR). CHOE, Tae-Boo [KR/KR]; Walkerhill Apt. 52-501, 145-8, Kwangjang-dong, Kwangjin-ku, Seoul 143-752 (KR). (74) Agent: KIM, Won-Ho; 702, Teheran Building, 825-33, Yoksam-dong, Kangnam-ku, Seoul 135-080 (KR).</p>		<p>(81) Designated States: CN, JP, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</p>

(54) Title: COMPOSITION, KIT, AND METHOD FOR DETECTING HELICOBACTER PYLORI IN BIOPSY

(57) Abstract

Disclosed are a composition for detecting Helicobacter pylori including urea 0.5 to 4 percent by volume, potassium phosphate 0.05 to 0.2 percent by volume, phenate reagent solution 0.8 to 1.7 percent by volume, an indicator having a pK_a of 6.5 to 8.5, 0.002 to 0.005 percent by volume, and a balance of water, a kit for detecting Helicobacter pylori using the composition and detecting method using the composition can determine quickly and accurately whether or not an infection by Helicobacter pylori exists, can obtain the same determination results after an elapse of time and can be used easily in an endoscope chamber.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COMPOSITION, KIT, AND METHOD FOR DETECTING HELICOBACTER PYLORI IN BIOPSIE

BACKGROUND OF THE INVENTION

5

(a) Field of the Invention

This invention relates to detection of *Helicobacter pylori* (hereinafter called *H. pylori*), source bacterium of gastrointestinal disorders, and in particular relates to an *H. pylori* detecting composition produced by applying a phenate
10 method for ammonium ion's quantitative determination to a conventional urease enzyme test and relates to a detecting kit and method using this composition.

(b) Description of the Related Art

H. pylori is noted as a source bacterium causing gastrointestinal
15 disorders including gastric cancer and gastroduodenal ulcers, and the WHO has prescribed *H. pylori* as a typed carcinogen. Precise diagnosis, i.e., accurate detection of *H. pylori*, should be promoted to cure gastrointestinal disorders caused by this *H. pylori*. *H. pylori* detecting methods being used up to now can be divided mainly into two types, i.e., non-invasive tests not using an
20 endoscope and invasive tests using an endoscope to get biopsy tissue for detection purposes.

Non-invasive tests include serology tests and ^{13}C or ^{14}C Breath tests. Serology tests measuring *H. pylori* antibodies have a problem in that the existence of the antibody does not necessarily mean that diseases are
25 developing because it takes more than 6 months for antibody levels to drop after the cure of diseases. The ^{13}C or ^{14}C Breath test uses the principle that if urea marked with ^{13}C or ^{14}C is ingested, it is transformed into marked CO_2 in the stomach if *H. pylori* exists and detected in exhaled air. However, the method is not usually used because it involves high cost equipment.

30 On the other hand, invasive tests are more commonly used to detect *H. pylori*. Invasive tests detecting methods using biopsy tissues samples collected during an endoscopy includes histology, the PCR (Polymerase

Chain Reaction) method, culturing, and the urease enzyme test. Histology is a method confirming the existence of *H. pylori* through general tissue research, the PCR method is a method confirming the existence of *H. pylori* by using chain reaction of polymerization enzyme, and culturing is a method of directly
5 cultivating *H. pylori* from biopsy tissue. However, there is a problem in that all these methods are difficult and take much time, so they can not be used commercially.

The urease enzyme test, which can be used easily in the endoscope chamber and is the most efficient method, uses the fact that *H. pylori* produces
10 urease enzyme which has a much higher degree of activity compared to other microorganisms. That is, if urease exists urea added in the detecting kit decomposes and ammonia is produced to cause pH to be increased making a pH indicator react and change color. A *H. pylori* detecting method measures the activity degree of urease like this. Commercialized kits using the urease
15 enzyme test now are "CLOtest" (USP No. 4,748,113), "Hpfast" (USP No. 5,439,801) and "Pyloritek" (USP No. 5,314,804 and 5,420,016). Compositions used in "CLOtest" include 10 ~ 40 g/l of urea, 1 ~ 5 g/l of bactericide, an available quantity of pK_a 6.5 ~ 8.5 indicator (phenol red) and the remainder being water, having a 5.0 ~ 6.5 pH. However, the CLOtest has problems in
20 that the positive rate vary with the readers because the reaction speed of urease that the above compositions and *H. pylori* produce is slow. Therefore, diagnosis is possible only after about 24 hours and it is not possible to obtain diagnostic results on the endoscope examination date which requires that the patient inconveniently visit the hospital once more. Furthermore, the various
25 colors appearing after more than 24 hours makes accurate diagnosis difficult. Hpfast is fundamentally similar to the CLOtest, but different in that cell wall detergent is added and uses as an indicator mixture of phenol red, methyl red, and bromothymol blue. With Hpfast, an *H. pylori* positive is determined from a dark green color of pH 6.2 and a light green color of pH 6.0 is determined as an
30 *H. pylori* negative. However, the accurate reading of this dark green and light green colors is difficult. Pyloritek uses a multilayer test device differently from

the above mentioned CLOtest and Hpfast, the main characteristics of this is that the urease reacting part and reacted product as an ammonia detecting part are on different layers and the pHs of each layers are different. Although Pyloritek can produce diagnosis results within one hour, the false positive rate increases if the determination time exceeds one hour. Therefore, there is an increasing possibility of false positives if an accurate reaction time is not adhered to during the endoscope examination.

Moreover, the above mentioned urease enzyme methods including the CLOtest, Pyloritek, etc. could also react with the small amounts of low activity urease which bacilli such as *Staphylococcus hominis*, *Streptococcus salivarius*, *E. aerofaciens*, *L. fermentum*, etc. have and false positive rates could be increased in cases of long decision times.

SUMMARY OF THE INVENTION

The present invention is designed to solve the above mentioned problems of the conventional technology by providing a *H. pylori* detecting composition, a *H. pylori* kit, and a method for using the above composition which allows the rapid and accurate determination of whether or not a *H. pylori* infection exists, which is the source bacterium that causes gastrointestinal disorders. The present invention also provides the same test results after a period of time passed, and it can easily be used in the chamber of an endoscope.

In order to achieve the above described purpose of this invention, this invention first provides *H. pylori* detecting composition including urea from 0.5 to 4 vol%, KH_2PO_4 from 0.05 to 0.2 vol%, phenate reagent solution from 0.8 to 1.7 vol%, an indicator from 0.002 to 0.005 vol% having a pKa of from 6.5 to 8.5 and a balance of water. Among the above described constituents, said phenate reagent solution from 0.8 to 1.7 vol% is preferably composed of manganous sulfate solution from 0.5 to 1 vol%, hypochlorite reagent from 0.2 to 0.5 vol% and phenate reagent from 0.1 to 0.2 vol % and the above described indicator is preferably phenol red. Moreover, said composition more preferably comprises

gelling agent from 0.5 to 2 vol%. Especially said composition most preferably
comprise 2 vol% urea, 0.05 vol% KH_2PO_4 , 1 vol% manganous sulfate solution,
0.5 vol% hypochlorite reagent, 0.2 vol% phenate reagent, 0.0025 vol% phenol
red, 1 vol% agar and a balance of water. And said composition preferably has
5 pH from 6.0 to 7.8.

Second, this invention is made from the above described composition
and provides an H. pylori detecting kit including biopsy tissue receptical test
device, positive control produced by further adding 10 ~ 20 μl of 0.1N NaOH
solution to said constituents and negative control made of said constituents
10 wherein no biopsy tissue is placed.

Third, this invention provides an H. pylori detecting method from biopsy
tissue including stages of reacting biopsy tissue with said composition and
observing color change of said composition.

15

BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying drawings, which are incorporated in and constitute a
part of the specification, illustrate an embodiment of the invention, and, together
with the description, serve to explain the principles of the invention:

Figure 1 is a perspective view of detecting kit according to this
20 invention;

Figure 2 is a sectional view of detecting kit according to this invention;
and

Figure 3 is a drawing describing application example of detecting kit
according to this invention.

25

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

In the following detailed description, only the preferred embodiment of
the invention has been shown and described, simply by way of illustration of the
best mode contemplated by the inventors of carrying out the invention. As will
30 be realized, the invention is capable of modification in various obvious respects,
all without departing from the invention. Accordingly, the drawings and


description are to be regarded as illustrative in nature, and not restrictive.

Hereinafter, further details of this invention are explained as follows:




In detecting H. pylori, the inventors tried to substantially lower false positive rates due to the subjectivity of the readers by transforming the existing urease enzyme tests from the methods conventionally used to a composition that uses a speedy change of colors that enables an accurate determination to be made rapidly by comparing the test results simultaneously with positive and negative controls. Namely, since test kits using existing urease enzyme tests detect the pH increase caused by OH⁻ ions produced from the reaction of water with the ammonia generated by simply decomposing urea, the inventors tried to eliminate the ammonium ions themselves in order to increase the rate of pH change. To do this, the inventors applied the phenate method (Standard Methods for the Examination of Water and Wastewater 19th ed. 1995, American Public Health Association, pp 4-80 ~ 4-82.) used in quantification of ammonium ions.

The Urea decomposition process by urease is presented as following reaction formula:

[reaction formula 1]




(urea)



The constituents of this present invention increase pH promptly by transforming the products of the above described urea decomposition process, i.e., ammonium ions, into indophenol blue to continue urea decomposition. In order to achieve the above described purpose, the phenate method, a remarkably effective method of measuring ammonia concentration in the aqueous solution, is applied. The phenate method, using the principle that ammonia under a manganic catalyst reacts with hypochlorite base and phenol

to change into indophenol blue, can be described as the following reaction formula:

[reaction formula 2]



That is, the constituents of this invention make the reaction speed faster in order to increase pH rapidly by adding phenate reagent solution to constituents used in the conventional urease enzyme test and transforming
10 ammonium ion generated in the urea decomposition process by the above described urease into indophenol blue.

Additionally, NaOCl is among the constituents that acts to reduce false positives that follow increased concentrations due to urease from bacilli as it
15 has an inhibiting effect on weak urease activity. This also allows test results to be obtained after longer testing time periods.

In this invention, the added quantity of phenate reagent solution is adjusted within a range so that H. pylori urease activity is not inhibited and color
20 change of the pH indicator is not affected. This invention's constituents and the possible content range of each components is as below:

urea from 0.5 to 4 vol%,
KH₂PO₄ from 0.05 to 0.2 vol%,
phenate reagent solution from 0.8 to 1.7 vol%,
25 0.002 ~ 0.005vol % of an indicator having pK_a from 6.5 to 8.5 and
a balance of water

From the above described composition, urea acts as a urease substrate which H. pylori produces, about 0.5 ~ 4 vol% of content is an appropriate level
30 to measure activity of urease which H. pylori produces. KH₂PO₄ acting as a buffer solution should include 0.05 ~ 0.2 vol%, i.e., it can not obtain desired the

pH range at lower than 0.05 vol% and tends to inhibit pH change at higher than 0.2 vol%. The phenate reagent solution, acting to transform ammonium ions into indophenol blue and to inhibit urease activities of other microorganisms should be included at 0.8 ~ 1.7 vol%, i.e., at lower than 0.8 vol% the effect of increasing the pH change speed by eliminating ammonium ions and the inhibiting effect of urease activity of other microorganisms are insufficient, while at higher than 1.7 vol% pH is increased much too high and the urease activities of other microorganisms as well as *H. pylori* tend to be inhibited. An indicator having a pKa from 6.5 ~ 8.5 acts to sense pH change, with phenol red being the most appropriate in this invention, preferably between 0.002 ~ 0.005 vol% for the precise determination. Phenol red has characteristics of showing yellow in an acid solution and a purplish-red color in a basic solution.

The above described phenate reagent solution containing in detail a manganous sulfate solution acting as a catalyst, hypochlorite reagent acting as a reactant reacted with ammonium ions and phenate reagent, and is especially desirably composed of 1 vol% manganous sulfate solution, 0.5 vol% hypochlorite reagent and 0.2 vol% phenate reagent. Moreover, it is further desirable that this constituent contains 0.5 ~ 2 vol% of a gelling agent, e.g., agar, as it can then be used conveniently in the form of a soft gel. It is preferable that constituent is controlled in the range of pH 6.0 ~ 7.8 because it aides precise determination when the pH is kept in this range.

Further desirable components and contents range of this invention's constituents are as follows:

2 vol% urea, 0.05 vol% KH_2PO_4 , 1 vol% manganous sulfate solution, 0.5 vol% hypochlorite reagent, 0.2 vol% phenate reagent, 0.0025 vol% phenol red, 1 vol% agar, and a balance of water.

On the other hand, desirable components and contents of stock solution of phenate reagent solution used in this invention are as follows:

Manganous sulfate solution; 0.05 vol% $\text{MnSO}_4 \cdot \text{H}_2\text{O}$, and a balance of distilled water,

Hypochlorite reagent; 1 vol% NaOCl , and a balance of distilled water,

Phenate reagent; 2.5 vol% NaOH, 8 vol% phenol, and a balance of distilled water.

This invention is to react with the above described constituents by placing biopsy tissue obtained through an endoscope on the gell made of the
5 above described constituents and observe color change. The gell is made of the above described constituents and a positive control gell is produced by adding to the above described constituents a minimum quantity of NaOH that can show positive results, i.e., 10 ~ 20 μ l 0.1N NaOH solution. A determination can be made as to whether or not there is an H. pylori infection
10 comparing the color of the negative control gell in which NaOH is not added at all. Accordingly, a test device with positive and negative controls placed side by side for comparison at a glance with colors indicating positive or negative results will decrease erroneous test results due to the subjectivity of the reader.

A perspective view of a detecting kit according to this invention is
15 presented in Figure 1 and side view in Figure 2, respectively. An application example of a detecting kit according to this invention is also presented in Figure 3. Referring to the above described Figures 1 to 3, 1 is a cover, 2 is a container, most desirably a transparent container made of an acrylic material, 3 is detecting composition according to this invention, 4 is a test device, 5 is the
20 biopsy specimen area, desirably opaquely treated such that biopsy specimen is not seen from the side, 6 is the negative control, 7 is the positive control and 8 is the biopsy sample.

A desirable practical example and comparative example of this invention are described. However, the below described practical example is
25 only one of practical examples of this invention and this invention is not limited to the below described practical example.

[Practical example 1]

Making H. pylori detecting compositions

30 First the below described composition of stock solution was prepared;
20 vol% urea, 2 vol% KH_2PO_4 , 0.01 vol% phenol red, and 2 vol% agar.

As stock solution of phenate reagent, 100 ml of manganous sulfate solution was made by adding distilled water to $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ 50 mg, 100 ml of hypochlorite reagent was made by adding distilled water to 10 ml of 10% NaOCl and the pH was adjusted to 6.8 with concentrated hypochloric acid.
5 100 ml of phenate reagent was prepared by adding distilled water to 2.5 g of NaOH and 8 ml of phenol. After the above described agar solution was autoclaved (121°C , 1.5 atm) for 15 minutes, it was left alone until it is used in the water tank at 55°C and the below described 2 X reagent were produced.

A 50 ml mixture was made by mixing the above described 10 ml urea
10 stock solution, 2.5ml KH_2PO_4 stock solution, 25 ml phenol red stock solution, 1 ml manganous sulfate solution, 0.5 ml hypochlorite reagent and 0.2 ml phenate reagent and adding distilled water. After bacilli of the above described produced 2 X reagent were filtered out by 0.2 μm filters, the same amount as 2% agar solution was mixed. As a result of that, the final concentration of the
15 constituents was as below and the final pH was 7.5 at this time:

2% urea, 0.05% KH_2PO_4 , 0.0005% manganous sulfate, 0.005% NaOCl , 0.2% phenate reagent (0.005% NaOH , 0.016% phenol), 0.0025% phenol red and 1% agar.

20

[Practical example 2]

Manufacture of a H. pylori detecting kit

The composition manufactured from example 1 was injected into one wall of a multiwall plate, test device with biopsy tissue obtained from an endoscope, a positive control was made by injecting into another wall the
25 composition manufactured and by further adding 10 μl of 0.1 N NaOH to the above described composition, a negative control was made by injecting the above described composition into another wall, thereby making an H. pylori detecting kit. The above described detecting kit is called "PET (Pylori Easy Test) kit".

30

[Test example 1]

After a patient's biopsy tissue was obtained, a CLOtest and PET of example 2 were used on that biopsy tissue simultaneously, and the positive rate per hour was compared and the results were described in the below Table

5 1.

[Table 1]

Positive rate per hour

Hour	PET	CLOtest
1	10	3
2	10 (+0)	3 (+0)
3	13 (+3)	8 (+5)
4	13 (+0)	8 (+0)
12	No test	No test
24	13 (+0)	10 (+2)
Total positive rate	13/26	10/26 or 17/26*

10 * 7 readings were difficult to determine.

As a result, PET could confirm 76.9% of total positives within one hour and 100 % after 3 hours. On the other hand, CLOtest could confirm only 17.6 ~ 30 % of the positives and 100% only after 24 hours. However, uncertain color which could not be distinguished distinctly appeared after 24 hours and
 15 the positive rate was varied from 38.4 % (10/26) to 65.3 % (17/26) depending on the observers. Therefore, we can see that rapid determination is possible in the case where PET is used according to this invention.

After that, we observed the degree of consistency of PET results with the PCR method. Here, the PCR method used an amplifying method of 26kDa
 20 protein gene, a method known as the most unique and sensitive in *H. pylori* detection (Ho, S, et al. 1991, Direct Polymerase Chain Reaction Test for detection of *H. pylori* in Human and Animals, J. Clin. Microbiol, 29 pp 2543 ~ 2549). The results were described the following table 2:

[Table 2]

Consistency ratio with PCR method

5

PET	CLOtest
26/26 (100 %)	16/26 (61.5 %): two false negatives, five false positives
	21/26 (80.7 %): when unreadable colors were judged to be negatives

From the above described table 2, we can see that all of the 26 biopsy tissues are consistent with the results of the PCR method when using this PET invention and that the consistency ratio ranges between 61.5 ~ 80.7% for the
 10 Clotest. Therefore, we can see that it is possible to judge that this PET invention is more precise than Clotest.

[Test example 2]

Biopsy tissue obtained during an endoscopy was refrigerated at -20
 15 °C for 1 week, and the PET and CLOtest were then performed according to the same method as in test example 1 with the results compared and presented in table 3 and table 4.

[Table 3]

20 Positive rate per hour

Hour	Practical example 1	CLOtest
1	6	1
2	11 (+5)	2 (+1)
3	13 (+2)	4 (+2)

4	13 (+0)	8 (+4)
12	13 (+0)	12 (+0)
24	13 (+0)	12 (+0)
Total positive rate	13/22	12/22*

* 6 readings were difficult to determine.

[Table 4]

Consistency with PCR

PET	CLOtest
22/22 (100 %)	21/22 (95.5%); one false negative
	18/22 (81.8 %); when unreadable colors were judged to be negatives

5

From table 3 and table 4, if PET according to this invention is used, it is notable that precise judgement is possible on biopsy tissue in which urease activity of *H. pylori* was decreased to certain degree.

This invention could provide *H. pylori* detecting compositions which tell promptly and precisely whether or not an *H. pylori* infection (the source bacterium causing gastrointestinal disorders) exists, and the same results can be obtained even after time has elapsed and the compositions are easily used in an endoscope chamber. The invention provides an *H. pylori* detecting kit and a method of using *H. pylori* detecting compositions.

15

[Test example 3]

Urease decomposition capacity was examined on *H. pylori* bacterium and *Staphylococcus hominis* bacterium separated from biopsy tissue obtained during an endoscopy in order to examine the *H. pylori* peculiarity of PET.

The examination method involved first having *H. pylori* bacterium and *S. hominis* bacterium cultivated and diluted in sterilized distilled water so as to have an appropriate number of bacilli, and then PET and CLOtest were performed and compared according to the same method as in test example 1.

5 The results were described in table 5.

[Table 5]

Peculiarity comparison on *H. pylori* bacterium and *S. hominis* bacterium

Classification		UBPH (0.02 %)	CLOtest
H. pylori	10 ⁵ cells	< 30 min	within 30 min
	10 ⁴ cells	1 h	2 h
	10 ³ cells	No response even after 24 hours	No response even after 24 hours
S. hominis	10 ⁷ cells	8 h	6 h
	10 ⁶ cells	No response even after 24 hours	10 h

10 From the above described table 5 it is noted that if PET according to this invention is used, high peculiarity detection is possible with urease activity of *H. pylori* showing a fast response and urease activity of other bacilli showing low response.

Although preferred embodiments of the present invention have been
 15 described in detail hereinabove, it should be clearly understood that many variations and/or modifications of the basic inventive concepts herein taught which may appear to those skilled in the present art will still fall within the spirit and scope of the present invention, as defined in the following claims.

WHAT IS CLAIMED IS:

1. A composition for detecting *Helicobacter pylori* including:
 - a) urea from 0.5 to 4 vol%;
 - b) KH_2PO_4 from 0.05 to 0.2 vol%;
 - 5 c) phenate reagent solution from 0.8 to 1.7 vol%;
 - d) an indicator having a pK_a of 6.5 to 8.5, 0.002 to 0.005 vol%; and
 - e) a balance of water.
2. The composition of claim 1 wherein said phenate reagent solution from 0.8 to 1.7 vol% of said *Helicobacter pylori* detecting composition comprises:
 - 10 a) manganous sulfate solution from 0.5 to 1 vol%;
 - b) hypochlorite reagent from 0.2 to 0.5 vol%; and
 - c) phenate reagent from 0.1 to 0.2 vol%.
3. The composition of claim 1 wherein said composition is a *Helicobacter pylori* detecting composition in which the indicator is phenol red.
- 15 4. The composition of claim 1 wherein said composition is a *Helicobacter pylori* detecting composition further including gelling agent from 0.5 to 2 vol%.
5. The composition of claim 4 wherein said composition is a *Helicobacter pylori* detecting composition including:
 - a) 2 vol% urea;
 - 20 b) 0.05 vol% KH_2PO_4 ;
 - c) 1 vol% manganous sulfate solution;
 - d) 0.5 vol% hypochlorite reagent;
 - e) 0.2 vol% phenate reagent;
 - f) 0.0025 vol% phenol red;
 - 25 g) 1 vol% agar; and
 - h) a balance of water
6. The composition of claim 1 wherein said composition is a *Helicobacter pylori* detecting composition having a pH from 6.0 to 7.8.
7. A kit for detecting *Helicobacter pylori* including:
 - 30 a) test device made of composition according to claim 4,
wherein biopsy tissue is placed;

- b) positive control manufactured by further adding 10 ~ 20 μ l of 0.1 N NaOH solution to composition of claim 4; and
- c) negative control manufactured from composition of claim 4, where biopsy tissue is not placed.

5 8. A method of detecting *Helicobacter pylori* from biopsy tissue comprising the steps of:

- a) reacting biopsy tissue and said composition according to claim 1; and
- b) observing the color change of the said biopsy tissue reacted composition.

1/2

FIG. 1






FIG. 2



2/2

FIG. 3



INTERNATIONAL SEARCH REPORT

International application No.
PCT/KR 99/00161

A. CLASSIFICATION OF SUBJECT MATTER

IPC⁶: C 12 Q 1/58; C 12 Q 1/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC⁶: C 12 Q 1/58; C 12 Q 1/04

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

WPI, CAS

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5 439 801 A (JACKSON), 08 August 1995 (08.08.95), column 2, lines 35-52; column 6, lines 17-21; column 7, lines 57,58; claims 1-6,9.	1,3-5,7,8
A	WO 97/30 351 A1 (THE BOOTS COMPANY PLC), 21 August 1997 (21.08.97), abstract; page 4, line 20-23; page 5, lines 2-4, 22,23; page 6, line 23- page 7, line 12; page 10, line 23 - page 11, line5.	1,2,5
A	US 5 420 016 A (BOGUSLASKI et al.), 30 May 1995 (30.05.95), column 6, lines 10-15; column 10, lines 5-12; claim 9.	1,3,7
A	EP 0 204 438 A2 (MARSHALL), 10 December 1986 (10.12.86), examples.	1,3-5,8

☐ Further documents are listed in the continuation of Box C.

☒ See patent family annex.

* Special categories of cited documents:

„A“ document defining the general state of the art which is not considered to be of particular relevance

„E“ earlier application or patent but published on or after the international filing date

„L“ document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

„O“ document referring to an oral disclosure, use, exhibition or other means

„P“ document published prior to the international filing date but later than the priority date claimed

„T“ later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

„X“ document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

„Y“ document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

„&“ document member of the same patent family

Date of the actual completion of the international search

25 June 1999 (25.06.99)

Date of mailing of the international search report

03 August 1999 (03.08.99)

Name and mailing address of the ISA/AT
Austrian Patent Office
Kohlmarkt 8-10; A-1014 Vienna
Facsimile No. 1/53424/200

Authorized officer

Mosser

Telephone No. 1/53424/437

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No.

PCT/KR 99/00161

In Recherchenbericht angeführtes Patentedokument Patent document cited in search report Document de brevet cité dans le rapport de recherche		Datum der Veröffentlichung Publication date Date de publication	Mitglied(er) der Patentfamilie Patent family member(s) Membre(s) de la famille de brevets	Datum der Veröffentlichung Publication date Date de publication	
US A	5439801	08-08-1995	CA AA WO A1	2160916 9521937	17-08-1995 17-08-1995
WO A1	9730351	21-08-1997	AU A1 GB A0	17696/97 9603031	02-09-1997 10-04-1996
US A	5420016	30-05-1995	AU A1 CA C EP A1 EP A4 JP T2 JP B2 US A WO A1	37361/93 2131317 633946 633946 7505279 2638682 5314804 9319200	21-10-1993 24-02-1998 18-01-1995 26-06-1996 15-06-1995 06-08-1997 24-05-1994 30-09-1993
EP A2	204438	10-12-1986	AT E BR A CA A1 DE CO DK A0 DK A EP A3 EP B1 HK A JP A2 JP B4 MX B NO A NO B NO C SG A US A AU A1 AU B2 ZA A	61414 8602243 1274757 3677820 2283/86 2283/86 204438 204438 351/92 62026000 6095960 165014 861966 170091 170091 148/92 4748113 57398/86 601363 8603605	15-03-1991 13-01-1987 02-10-1990 11-04-1991 16-05-1986 18-11-1986 27-05-1987 06-03-1991 22-05-1992 03-02-1987 30-11-1994 11-08-1993 18-11-1986 01-06-1992 09-09-1992 29-01-1993 31-05-1988 20-11-1986 13-09-1990 27-01-1988